

**HOW TO STOP OBSESSIVE WORRY:
A KEY TO PEACE OF MIND
William Doverspike, Ph.D.
Drdoverspike.com
770-913-0506**

This article is designed to be educational in nature and it is not intended to provide professional advice. The reader is encouraged to contact a licensed mental health professional if professional advice, diagnostic consultation, or treatment is being sought.

There is a story of an old tombstone that contained the inscription: “95% of what put me here never happened.” Yet most people worry about things occasionally, which can be adaptive and useful if it motivates the person into taking constructive action such as preparing for a test, considering a job change, or planning for the future. However, when worrying becomes excessive, unproductive, or difficult to control, it may be maladaptive or even pathological. Difficulty controlling worrying is one of the primary features that distinguishes pathological worry from the everyday anxieties that most people experience when they anticipate and prepare for an important event. Pathological worrying is sometimes described as obsessive worrying, which may become a true obsession (from the Latin *obsessus*, meaning “besieged”). The Latin root is instructive because when a person is obsessed, it is as if the person were besieged by uncontrollable thoughts.

Obsessions are intrusive thoughts that a person tries to control, resist, or eliminate. In other words, an obsession is a recurring thought that outweighs any other thought. There are some important differences between true obsessions and worry. For example, worry involves an apprehensive expectation about some type of realistic domain of activities or events (e.g., family, health, finances, or school/work), whereas true obsessions have an irrational quality (e.g., a contamination phobia). In general, true obsessions are not as common as obsessive worry.

In order to cope with the weight of obsessive worry, psychologists have identified several cognitive-behavioral activities that can be useful in reducing the frequency and intensity of obsessive thoughts. Some of these same ideas are contained in a variety of sources such as *The Anxiety and Worry Workbook* (Clark & Beck, 2012), *The Anxiety and Phobia Workbook* (Bourne, 2010), and other sources.

Obsessive worry is like a negative spiral. The longer you spend time with an obsession, the deeper into it you can get stuck. Neurologically, through the process of synaptic neuroplasticity, the repetition or practice of an activity or thought reinforces neural pathways much like a well trod path in a forest. Psychologically, a mental obsession can also be viewed as a form of trance (Nakken, 1996). The more you induce it by repetition, the more entranced you become and the more difficult it may be to “break the spell.”

The downward pull of an obsessive spiral can be very compelling. Following the path of least resistance is likely to keep you going round and round until you are eventually feeling down and out. Deliberately choosing to break out of the obsessive thinking may be difficult at first, especially if you’re highly anxious. With practice, it can become easier. Below are some examples of alternative activities and experiences that will help you shift your mind away from obsessive thinking.

A deliberate act of will is required to stop obsessing. You need to make a deliberate effort to move away from circular mental activity and get out of your head by “shifting gears” to another modality of experience, such as bodily activity, expressing emotions, interpersonal communication, sensory

distraction, or a specific ritual. In some cases, an alternative, positive obsession may help. For example, the vicious cycle of negative worry can sometimes be replaced by positive obsessions such as uplifting thoughts, affirmations, or slogans (e.g., “This too shall pass”).

Find an alternative positive obsession. In addition to focusing on uplifting affirmations or slogans, there are other types of “positive obsessions” that can be helpful. For example, try performing a complicated mental activity, solving a complex puzzle, practicing a mental agility game, studying an interesting subject, or even writing letters or articles.

Engage in positive physical activities. Although difficult at first, engaging in physical activities can help break an obsessive cycle. Physical activities can include engaging indoor or outdoor exercise, performing household chores, or simply taking a walk. The more entrenched the obsessive cycle, the more vigorous the activity may be needed to dig out of the rut.

Use visual and sensory-motor distractions. Find an enjoyable activity in which you can “lose yourself” in the moment of the activity itself. Active distractions are usually more effective than passive ones (e.g., watching a movie). Effective distractions can involve working on your computer, playing a complex video game, engaging in uplifting reading, or even digging in the dirt. Spend time engaging in arts and crafts, repairing something, or gardening. Any healthy activity can be useful if it is one in which you lose track of time while engaging in the experience of the activity.

Listen to evocative music to release repressed feelings. Repressed feelings—particularly dysphoric emotions such as sadness or anger—may underlie and “drive” the obsessive thinking. In such cases, music may help you get in touch with your feelings that are below the surface of your

obsessions. This relationship may be the basis of the adage that “music soothes the savage soul.”

Talk to someone. Use the old adage, “Don’t worry alone.” Instead, talk with a trusted person about something other than the worry, unless you want to express your feelings about it. People who engage in obsessive thinking tend to focus on their thoughts rather than their feelings. If you can learn to share your *feelings*, it may help dissipate your obsessive thinking.

Use visual imagery. Some research suggests that chronic worriers are people who engage in frantic, intense thought processes without accompanying images of the threat (Borkovec, Alcaine, & Behar, 2004; Borkovec, Shadick, & Hopkins, 1991; Fisher & Wells, 2009; Roemer & Borkovec, 1993). In other words, chronic worriers may spend so much time thinking about upcoming problems that they don’t have any attentional activity left for the important process of creating visual images of the potential threat (Craske & Barlow, 2006). With the help of a psychologist, you may be able to learn how to form mental images associated with negative affect which, paradoxically, can be processed in a way that helps reduce anxiety and worrying.

Consider using positive imagination. Whereas creativity involves the positive use of imagination, worry involves the negative use of imagination. People who worry often imagine the worst case scenario of what might happen. However, the best case scenario is often as likely as the worst case scenario. Rather than thinking of the worst short term consequence of what might happen, use imagination to visualize the best long-term consequence of what might happen.

Use progressive muscle relaxation. Progressive muscle relaxation is a stress management technique by which you can learn to reduce anxiety by learning how to relax the muscle tension. Relaxation exercises can be done alone or in

combination with abdominal breathing, which means breathing fully from your abdomen or from the bottom of your lungs. It is the reverse of the way you breathe when you're anxious or tense, which typically involves breathing that is shallow and high in your chest. Practice progressive relaxation for 5-10 minutes until you feel fully relaxed and free from obsessive thoughts.

Consider thought-stopping techniques. Based on Cautela and Wisocki's (1977) original article, behaviorists sometimes use cognitive interventions designed to reduce unwanted thoughts. One common technique is to snap a rubber band placed on the wrist with the goal of providing an aversive consequence whenever the unwanted thought occurs. However, some psychologists (e.g., Leahy, 2006, 2010) believe thought stopping not only doesn't work but that it can lead to *thought-rebounding*, which refers to an increase in unwanted thoughts that makes matters worse. For example, suppressing thoughts of white bears can lead to an increase or rebound of these thoughts after the suppression (Wegner, 1989). The purpose of the rubber band is to increase one's self-awareness. Rather than stopping a thought, it may be more helpful to redirect one's thoughts (i.e., Stop, Interrupt, and Redirect).

Reframe worry as being less important. Rather than thinking of obsessive worry as something that hijacks your thoughts, think of it simply as a heckler wanting to be noticed. Just as worry can be empowered by giving it too much attention, it can also be disempowered as a heckler by putting it on the back row rather than in the front of the audience.

Practice healthy rituals. Combine abdominal breathing with a positive affirmation that has personal significance. Keep this up for 5-10 minutes, or until you're fully relaxed. This type of activity can function as a positive trance induction that can overcome the negative trance enforced by the obsessive worry. For example, an affirmation

can involve a statement such as "These are just thoughts" or simply "Let it go."

Consider meditation procedures. Although many of the above procedures focus on avoiding obsessive thoughts, psychologists have also developed some meditational approaches that focus on acceptance rather than avoidance of distressing thoughts and feelings (Roemer, Orsillo, & Barlow, 2002). Meditational procedures help people learn how to be more tolerant of these feelings which, contrary to common sense, can sometimes help a person reduce their obsessive thoughts.

Consider stopping thought meditation. The concept of *stopping thought* refers to the Zen mental state of samādi. In this state of meditative consciousness, normal mental chatter slows down or even stops for a period of time, which allows one to experience peace and serenity. Samādi is not achieved through force of will or normal intent, but rather through a combination of acceptance and returning to or focusing on a familiar state. For example, practitioners of zazen meditation typically sit while suspending all judgmental thinking and by letting ideas, images, and thoughts pass by without getting involved in them. It is a way of "letting go" of thoughts.

When all else fails, consider using prayer. In the words of Abraham Lincoln, "I have been driven many times upon my knees by the overwhelming conviction that I had nowhere else to go. My own wisdom and that of all about me seemed insufficient for that day."

Consider using the Serenity Prayer. From a spiritual perspective, excessive worry can be viewed as the antithesis of prayer. To counteract the toxic effects of an obsessive cycle, a healthy antidote can be found in the wisdom of theologian Reinhold Niebuhr's (1892–1971) Serenity Prayer, which was first popularized in the Alcoholics Anonymous *Grapevine* (1950, pp. 6-7):

God,
Grant me the serenity
to accept the things I cannot change,
the courage
to change the things I can,
and the wisdom
to know the difference.

Be realistic rather than idealistic. Despite the catchy title of this article, an accurate description of a realistic goal would be to manage obsessive worry rather than expecting to be completely free of worry. Again, some worry can be adaptive—when it leads to constructive action—and a realistic goal would be to transform worry into useful actions.

Consider medical consultation. Although beyond the scope of this article, there are medications that have been shown to be useful in the treatment of anxiety conditions. However, it is important to remember that “pills are not skills.” If regular practice of the above skills does not seem to help control your obsessive worries, it may be helpful to arrange a consultation with a board certified psychiatric physician concerning the benefits and risks of psychotropic medications.

Resources

Spitalnick, J., and Munford, M. (2021). *Raising resilience: 25 tips for parenting your child with anxiety or OCD*. Atlanta, GA: Anxiety Specialists of Atlanta.

Josh Spitalnick, PhD, ABPP is a Licensed Psychologist who is Board Certified in Behavioral and Cognitive Psychology by the American Board of Professional Psychology (ABPP). Marti Munford, LPC is a Licensed Professional Counselor.

The Spitalnick and Munford (2021) article is available at this [link](#).

Permission to upload and link was provided by the first author, Josh Spitalnick, PhD, ABPP at 12:11 PM on 05-26-2023.

References

- Alcoholics Anonymous. (1950). *Grapevine*. New York, NY: Alcoholics Anonymous World Service Inc.
- Borkovec, T. D., Alcaine, O. M., Behar, E. (2004). Avoidance theory of worry and generalized anxiety disorder. In R. G. Heimberg, C. L. Turk, & D. S. Mennin (Eds.), *Generalized anxiety disorder: Advances in research and practice* (pp. 77–108). New York, NY: Guilford.
- Borkovec, T. D., Shadick, R., & Hopkins, M. (1991). The nature of normal and pathological worry. In R. M. Rapee & D. H. Barlow (Eds.), *Chronic anxiety, generalized anxiety disorder, and mixed anxiety depression*. New York, NY: Guilford.
- Bourne, Edmund, J. (2010). *The anxiety & phobia workbook* (5th ed.). Oakland, CA: New Harbinger Publications, Inc. Pages 425–426.
- Clark, D.A., & Beck, A. T. (2012). *The anxiety & worry workbook: The cognitive-behavioral solution*. New York, NY: Guilford Press.
- Cautela, J. R., & Wisocki, P. A. (1977). The thought-stopping procedure: Description, application, and learning theory interpretations. *Psychological Record* 27,(1), 255–264.
- Craske, M. G., & Barlow, D. H. (2006). *Mastery of your anxiety and worry*. New York, NY: Oxford University Press.
- Fisher, P. L., & Wells, A. (2009). Psychological models of worry and generalized anxiety disorder. In M. M. Antony & M. B. Stein (Eds.). *Oxford handbook of anxiety and related disorders* (pp. 225–237). New York, NY: Oxford University Press.
- Leahy, R. L. (2006). *The worry cure: Seven steps to stop worry from stopping you*. New York, NY: Three Rivers Press.
- Leahy, R. L. (2010). Why thought stopping doesn't work. *Psychology Today*. <https://www.psychologytoday.com/blog/anxiety-files/201007/why-thought-stopping-doesn-t-work>
- Nakken, C. (1996). *The addictive personality: Understanding the addictive process and compulsive behavior* (2nd ed.). Center City, MN: Hazelden.
- Roemer, L., & Borkovec, T. D. (1993). Worry: Unwanted cognitive activity that controls unwanted somatic experience. In D. M. Wegner & J. W. Pennebaker (Eds.). *Handbook of mental control*. Englewood Cliffs, NJ: Prentice Hall.
- Roemer, L., Orsillo, S. M., & Barlow, D. H. (2002). Generalized anxiety disorder. In D. H. Barlow, *Anxiety and its disorders: The nature and treatment of anxiety and panic* (2nd ed.). New York, NY: Guilford Press.
- Wegner, D. M. (1989). *White bears and other unwanted thoughts: Suppression, obsession, and the psychology of mental control*. New York, NY: Penguin.

Correct Citation for Reference Entry

The reference entry correct citation styles for this document are illustrated below. Students should defer to the style preferences of their individual course instructors to determine whether the course instructor has preferences that are more specific than those shown below:

American Psychological Association

Doverspike, W. F. (2008). How to stop obsessive worry.
<http://drwilliamdoverspike.com/>

Chicago Manual of Style / Kate Turabian

Doverspike, William, "How to stop obsessive worry," 2008.
<http://drwilliamdoverspike.com/>

Note: According to the Chicago Manual of Style, blog posts are typically not included in bibliographies, but can be cited in the running text and/or notes. However, if a blog is cited frequently, you may include it in the bibliography.

Modern Language Association

Doverspike, William F. "How to stop obsessive worry" 2008
 [Date accessed]

Note: MLA guidelines assume that readers can track down most online sources by entering the author, title, or other identifying information in a search engine or a database. Consequently, MLA does not require a URL in citations for online sources such as websites. However, some instructors still ask for it, so check with your instructor to determine his or her preference.

Documentation

This document is cross-referenced to a portable document file (PDF) published from this Word document file: How to Stop Obsessive Worry.doc

Server path:

http://drwilliamdoverspike.com/files/how_to_stop_obsessive_worry.pdf

Server file name:

how_to_stop_obsessive_worry.pdf

Website tab: Practices [Popular Articles]

Link name: How to Stop Obsessive Worry

Workshop Presentation Deck:

N/A

Webinar Presentation Deck:

N/A

Workshop Handouts:

N/A

Published 2008 | Last updated 2024

Copyright © 2008 by William F. Doverspike, Ph.D.

Citations last updated 2024

The correct citation for this article is Doverspike, W. F. (2008, December). How to stop obsessive worry: A key to peace of mind. <http://drwilliamdoverspike.com/>